



## Automata and Logic

### Exercise Sheet 5

Dr. rer. nat. Rafael Peñaloza  
Summer Semester 2013

#### Exercise 27

Let  $V$  be the class of all finite semigroups  $S$  such that for all idempotent elements  $e \in S$ , we have  $Se = e$ . Show that  $V$  is an S-variety ultimately defined by

$$yx^n = x^n \quad (n \geq 1).$$

#### Exercise 28

Let  $\Sigma := \{a, b, c, d\}$ .

a) For  $L \subseteq \Sigma^*$  with

$$L := \{w \in \Sigma^* \mid w \text{ starts with } a \text{ or } b\} \cap \\ \{w \in \Sigma^* \mid |w| \geq 3 \text{ and } w \text{ starts and ends with the same symbol}\},$$

give a quantifier-free formula  $\phi$  using the signature  $\{Q_a, Q_b, Q_c, Q_d, <, \min, \max, s, p\}$  such that  $L(\phi) = L$ .

b) Let

$$\phi := \neg(\neg Q_a(s(s(p(s(\min)))))) \vee (s(\min) < p(p(\max))).$$

Use the method described in the proof of Prop. 2.11 to describe  $L(\phi)$  as a Boolean combination of languages from the set  $\{u\Sigma^* \mid u \in \Sigma^*\} \cup \{\Sigma^*u \mid u \in \Sigma^*\}$ .

#### Exercise 29

Let  $\Sigma$  be an alphabet. A language  $L \subseteq \Sigma^*$  is called *definite* for  $\Sigma$  if there exists an  $n \in \mathbb{N}$  such that we have for all  $w \in L$ :

$$\text{if } w = uv \text{ with } |u| = n, \text{ then } u\Sigma^* \subseteq L.$$

Show that  $L \subseteq \Sigma^*$  is definite for  $\Sigma$  iff  $L$  is a Boolean combination of languages of the form  $\{w\Sigma^* \mid w \in \Sigma^*\}$ .

### Exercise 30

Let  $\Sigma = \{0, 1\}^k$ . Show that the following is equivalent:

- $L$  is definite for  $\Sigma$ .
- There exists a quantifier-free closed first-order formula  $\phi$  over the signature  $\{P_1, \dots, P_k, <, \min, s\}$  with  $L(\phi) = L \setminus \{\varepsilon\}$ .

### Exercise 31

Let  $\Sigma, \Gamma$  be two alphabets, and let  $L \subseteq \Sigma^*$ . Prove or refute the following:

- $L \in \text{SF}_\Sigma \implies L \in \text{SF}_{\Sigma \cup \Gamma}$
- $L \in \text{SF}_{\Sigma \cup \Gamma} \implies L \in \text{SF}_\Sigma$

### Exercise 32

For  $\Sigma = \{a, b\}$ , check whether the following languages are star-free:

- $L_1 = (ab)^*$
- $L_2 = \{w \mid |w|_a = 3k \text{ for some } k \in \mathbb{N}\}$
- $L_3 = a(aba)^*b$

Use Thm. 3.6 from the lecture or give a star-free description of the language.