



Automata and Logic

Exercise Sheet 8

Dr. rer. nat. Rafael Peñaloza
Summer Semester 2013

Exercise 42


Let Σ be an alphabet. Prove the following:

- If $L \subseteq \Sigma^+$ is regular, then there exists a non-deterministic finite automaton \mathcal{A} with only *one* final state such that $L = L(\mathcal{A})$.
- If $L \subseteq \Sigma^*$ is regular, then there exists a non-deterministic finite automaton \mathcal{A} with at most *two* final states such that $L = L(\mathcal{A})$.
- There is *no* $k \geq 1$ such that the following holds:
If $L \subseteq \Sigma^\omega$ is Büchi recognisable, then there exists a Büchi automaton \mathcal{A} with at most k final states such that $L = L_\omega(\mathcal{A})$.

Hint: Consider the languages $a^\omega \cup b^\omega, a^\omega \cup b^\omega \cup c^\omega, \dots$

Exercise 43

Consider Büchi automata using the following transition system:



Check whether the recognised ω -language is empty for the following sets of final states:

- $F = \{q_0, q_1\}$
- $F = \{q_2, q_3\}$
- $F = \{q_1, q_3\}$

Exercise 44

For a finite automaton \mathcal{A} , let \mathcal{A}_{det} denote the minimal deterministic finite automaton such that $L(\mathcal{A}) = L(\mathcal{A}_{\text{det}})$. Prove or refute the following:

- a) $\lim L(\mathcal{A}) = L_\omega(\mathcal{A}_{\text{det}})$
- b) $L_\omega(\mathcal{A}) \subseteq L_\omega(\mathcal{A}_{\text{det}})$
- c) $L_\omega(\mathcal{A}_{\text{det}}) \subseteq L_\omega(\mathcal{A})$

Exercise 45

Let $(r_n)_{n \geq 0}$ be a sequence of real numbers. Show that there exists an infinite sub-sequence of $(r_n)_{n \geq 0}$ that is

- strictly increasing, or
- strictly decreasing, or
- constant.