



## Term Rewriting Systems

### Exercise Sheet 1

Prof. Dr.-Ing. Franz Baader / Dr. rer. nat. Marcel Lippmann  
Summer Semester 2016

#### Exercise 1

Consider the reduction system  $(M, \rightarrow)$  with  $M = \{A_1, A_2, A_3, A_4, B_1, B_2, B_3, C_1, C_2, C_3, C_4, D, E\}$  and  $\rightarrow \subseteq M \times M$ :

- $A_1 \rightarrow B_1, A_1 \rightarrow B_2, A_2 \rightarrow B_1, A_2 \rightarrow B_2, A_3 \rightarrow B_3, A_4 \rightarrow B_3,$
- $B_1 \rightarrow C_1, B_2 \rightarrow C_2, B_2 \rightarrow C_3, B_3 \rightarrow C_1, B_3 \rightarrow C_2, B_3 \rightarrow C_3, B_3 \rightarrow C_4,$
- $C_3 \rightarrow E, C_4 \rightarrow E,$  and
- $D \rightarrow C_4.$

Answer the following questions.

a) Which of the following properties are satisfied by  $\rightarrow$ ? Justify your answer.

- |                    |                |
|--------------------|----------------|
| i) finite          | iv) reflexive  |
| ii) symmetric      | v) irreflexive |
| iii) antisymmetric | vi) transitive |

b) Describe the following *closures*:

$\xrightarrow{=}, \xrightarrow{+}, \xrightarrow{*},$  and  $\leftrightarrow$ .

#### Exercise 2

Let  $\rightarrow$  be the *symbolic differentiation relation* introduced in the lecture.

a) Compute the *normal forms* of the following terms:

- $D_X(((X * X) * X) + (X * X)),$  and
- $D_X((X * Y) + (Y * Y)).$

b) Prove that  $\rightarrow$  is *terminating*.

#### Exercise 3

In the lecture, a *group* was defined by the following identities:

$$(x \circ y) \circ z \approx x \circ (y \circ z) \quad (\text{G1})$$

$$e \circ x \approx x \quad (\text{G2})$$

$$i(x) \circ x \approx e \quad (\text{G3})$$

a) Prove that groups satisfy the property that  $e$  is a right unit, i.e.

$$x \circ e \approx x \tag{G2'}$$

by showing that  $x \circ e$  can be transformed to  $x$  using the identities G1, G2 and G3.

b) Consider the following identity:

$$x \circ i(x) \approx e \tag{G3'}$$

Prove that G1, G2 and G3' do not imply G2'.

**Hint:**

Find a model of G1, G2 and G3' such that G2' does not hold in this model; such a model exists with only two elements.

#### Exercise 4

Consider the following identities:

$$(x \circ y) \circ z \approx x \circ (y \circ z) \tag{R1}$$

$$(x \circ y) \circ x \approx x \tag{R2}$$

Prove or refute whether the following identities are implied by R1 and R2.

a)  $(x \circ x) \approx x$

b)  $(x \circ y) \circ z \approx x \circ z$