**Lehrstuhl für Automatentheorie**
Institut für Theoretische Informatik, TU Dresden
Prof. Dr. F. Baader

Hans-Grundig-Str. 25
Dresden Johannstadt
Tel.: 0351/463–38237
Tel.: 0351/463–39171

# 8. Exercises for the Course „Automata and Logic“

**Exercise 36:**

Let $\Sigma = \{a\}$. As in the lecture, let $L_{k,n}$ denote the set of all first-order formulas (over the non-logical symbols $=$, $<$, and $Q_a$) containing $k$ free variables and having quantifier depth at most $n$. For the following combinations of $k$ and $n$, determine a *finite* set $\Gamma_{k,n}$ such that, for each formula $\varphi \in L_{k,n}$, there exists a $\psi \in \Gamma_{k,n}$ with $\varphi \equiv \psi$.

(a) $k = 1, n = 0$;

(b) $k = 2, n = 0$;

(c) $k = 0, n = 1$;

(d) $k = 1, n = 1$;

**Exercise 37:**

For the combinations of $k$ and $n$ listed in the previous exercise, determine the equivalence classes of the relations $\equiv_{k,n}$. To do this, use the sets $\Gamma_{k,n}$ determined in the previous exercise.

**Exercise 38:**

Using the results from the previous exercise, give the formulas $\phi_W$ for each equivalence class $W$ of $\equiv_{2,0}$. Then determine a finite disjunction of formulas $\phi_W$ for $\equiv_{k,n}$-classes $W$ which is equivalent to the following formulas:

(a) true

(b) $\neg(x < y) \vee x = y$

(c) false