

11. Exercises for the Course „Logic-based Knowledge Representation“


Exercise 35:

In the lecture, the translation of \mathcal{EL} -concepts into description trees has been presented on an intuitive level, only. Define this translation formally using induction on the structure of \mathcal{EL} -concepts.


Exercise 36:

Compute all possible homomorphisms between the following description trees:

1.



2.



Exercise 37:

Let \mathcal{L} be a Description Logic and C , D , and E be \mathcal{L} -concept terms. Prove that the following properties hold:

- (a) $\text{lcs}(C, D) = \text{lcs}(D, C)$
- (b) $\text{lcs}(C, \text{lcs}(D, E)) = \text{lcs}(\text{lcs}(C, D), E)$
- (c) The least common subsumer of C and D is unique up to equivalence.

Exercise 38:

Compute the least common subsumer of

(a) the following pairs of \mathcal{FL}_0 -concept terms:

$$C = A_1 \sqcap A_2 \sqcap \forall r.(A_1 \sqcap \forall s.A_1 \sqcap \forall r.A_2) \sqcap \forall r.\forall s.\forall r.(A_1 \sqcap A_2 \sqcap \forall s.A_2)$$

$$D = A_2 \sqcap \forall r.\forall r.(A_2 \sqcap A_3) \sqcap \forall s.A_2 \sqcap \forall r.(A_2 \sqcap \forall s.(A_2 \sqcap \forall r.(A_2 \sqcap A_3)))$$

(b) the following pairs of \mathcal{EL} -concept terms:

$$C = A \sqcap \exists r.(A \sqcap \exists(A \sqcap B)) \sqcap \exists r.(B \sqcap \exists s.B) \sqcap \exists s.(A \sqcap \exists r.\top)$$

$$D = B \sqcap \exists r.(A \sqcap B \sqcap \exists s.(A \sqcap B \sqcap \exists s.(A \sqcap B))) \sqcap \exists s.(A \sqcap B) \sqcap \exists s.(B \sqcap \exists r.\top)$$

Exercise 39:

Consider the following \mathcal{EL} -concept terms:

$$C_1 = \exists r.(A \sqcap \exists r.(A \sqcap B \sqcap \exists r.(A \sqcap B))) \sqcap \exists r.(B \sqcap \exists r.(A \sqcap B \sqcap \exists r.(A \sqcap B)))$$

$$C_2 = \exists r.(A \sqcap B \sqcap \exists r.(A \sqcap \exists r.(A \sqcap B)) \sqcap \exists r.(B \sqcap \exists r.(A \sqcap B)))$$

$$C_3 = \exists r.(A \sqcap B \sqcap \exists r.(A \sqcap B \sqcap \exists r.A \sqcap \exists r.B))$$

Compute $D = \text{lcs}(C_1, \text{lcs}(C_2, C_3))$ and prove that there is no \mathcal{EL} -concept term E such that D and E are equivalent and E 's length is smaller than $2^4 - 1$ (i.e., the number of nodes in a tree of depth 3).