



## Term Rewriting Systems

### Exercise Sheet 4

Prof. Dr.-Ing. Franz Baader  
Winter Semester 2011/2012

#### Exercise 15

The reduction relation  $\rightarrow$  enjoys the *diamond property* if

$$y_1 \leftarrow x \rightarrow y_2 \implies \exists z. y_1 \rightarrow z \leftarrow y_2.$$

Prove that, if  $\rightarrow$  enjoys the diamond property, then every element  $x$  is either in normal form or does not have a normal form.

#### Exercise 16

Let  $(A, \rightarrow_1 \cup \rightarrow_2)$  be the reduction system obtained from the reduction systems  $(A, \rightarrow_1)$  and  $(A, \rightarrow_2)$  by building the union of the two reduction relations.

Prove or refute: If  $\rightarrow_1$  and  $\rightarrow_2$  are confluent, then so is  $\rightarrow_1 \cup \rightarrow_2$ .

#### Exercise 17

Does *strong confluence* imply the following property?

$$y_1 \leftarrow x \rightarrow y_2 \implies \exists z. y_1 \xrightarrow{\equiv} z \xleftarrow{\equiv} y_2$$

Give a proof or counterexample.

#### Exercise 18

Consider the terms  $s = f(x, g(h(k(k(y))), x), z), h(x, y))$  and  $t = g(z, h(x, k(k(y))))$ .

Describe  $t|_1$ ,  $t|_{1111}$ ,  $t|_{11111}$ ,  $t[s]_2$ , and  $t[s]_2|_{21}$ .

#### Exercise 19

Prove the second part of Lemma 3.4 by induction on the length of words denoting positions:

If  $p \in \text{Pos}(s)$  and  $q \in \text{Pos}(t)$ , then

$$\begin{aligned} (s[t]_p)|_{pq} &= t|_q \\ (s[t]_p)[r]_{pq} &= s[t[r]_q]_p \end{aligned}$$

#### Exercise 20

Let  $E$  be a set of identities and  $\rightarrow_E$  the induced reduction relation. Prove that  $\rightarrow_E$  and  $\xrightarrow{*}_E$  are rewrite relations, i.e. are closed under substitutions and compatible with  $\Sigma$ -operations.