



Description Logics

Exercise Sheet 12

Dr.-Ing. Anni-Yasmin Turhan
Winter Semester 2012/13

Exercise 37

Let \mathcal{T} be a general \mathcal{EL} -TBox and $\hat{\mathcal{T}}$ the TBox obtained from exhaustive application of the normalisation rules NF1–NF5 to \mathcal{T} .

Finish the proof of Lemma 5.18 by showing the following:

- $\hat{\mathcal{T}}$ can be obtained from \mathcal{T} in polynomial time.
- $\hat{\mathcal{T}}$ is in normal form.
- For all concept names A, B occurring in \mathcal{T} , we have $A \sqsubseteq_{\mathcal{T}} B$ iff $A \sqsubseteq_{\hat{\mathcal{T}}} B$.

Exercise 38

Consider the TBox \mathcal{T} consisting of the following axioms:

$$\begin{aligned} A &\sqsubseteq B \sqcap \exists r.C \\ B \sqcap \exists r.B &\sqsubseteq C \sqcap D \\ C &\sqsubseteq \exists r.A \sqcap B \\ \exists r.\exists r.B \sqcap D &\sqsubseteq \exists r.(A \sqcap B) \end{aligned}$$

Check whether the following subsumption relations follow:

- $A \sqsubseteq B$
- $A \sqsubseteq \exists r.\exists r.A$
- $B \sqcap \exists r.A \sqsubseteq \exists r.C$